

# TORSIONAL SPLITTING AND FOUR-FOLD BARRIER TO INTERNAL ROTATION: THE ROTATIONAL SPECTRA OF VINYLSULFUR PENTAFLUORIDE

W. ORELLANA, SUSANNA L. STEPHENS, WALLACE C. PRINGLE, STEWART E. NOVICK, *Department of Chemistry, Wesleyan University, Middletown, CT, USA*; PETER GRONER, *Department of Chemistry, University of Missouri - Kansas City, Kansas City, MO, USA*; S. A. COOKE, *Natural and Social Science, Purchase College SUNY, Purchase, NY, USA*.

The rotational spectra of vinylsulfur pentafluoride, and three isotopologues (S-34 and both C-13's) have been recorded in the frequency region of 6 GHz to 20 GHz. Measurements were made using both cavity and chirped pulse Fourier transform microwave spectrometers. The four-fold barrier to the internal rotation of the  $-\text{SF}_5$  group against the vinyl group has been approximated from the spectral data which is possible due to the observation of easily resolved pure rotational transitions in each of the  $A$ ,  $B$ , and doubly degenerate  $E$  torsional substates. All transitions were successfully fit simultaneously using the ERHAM code. We note that this work, we believe, represents the first use of pure rotational spectroscopy to characterize a four-fold barrier internal rotation problem. Rotational constants, structure, and the internal rotation barrier height will be presented and compared to results from quantum chemical calculations.

